

Reverse Shoulder Replacement for Fracture

Dr. McCormick

What to Expect After Reverse Shoulder Replacement for Shoulder Fracture

Restrictions

For the first 6 weeks you may use your hand for writing and typing

You may drive when out of sling, usually at 6 weeks post surgery

you must be able to react quickly so try it in an empty parking lot if unsure.

Surgical site care

keep your initial dressing dry

it is normal to notice some drainage/discoloration of the dressing

nurses at the hospital will change your dressing before you are discharged

you may shower with the island dressings on but do not scrub the area

Pain management

Rest wear the sling until the block wears off

remove the sling 3 times per day to:

move your hand, wrist and elbow to keep these from getting stiff

do shoulder pendulum exercises

the sutures holding your repaired tendons are not strong enough to withstand the force of normal use

Ice 30 minutes per hour – remember to keep something between the cold pack/ice and your skin to avoid frostbite

Take pain medications as prescribed. If you have a reaction to the medications then call the office at 715.858.4650

Follow-up clinic visit

usually at 2 weeks after surgery

we will check your surgical site, review Xrays and answer any questions you may have

When to call the office

reactions to medications

dressing discoloration spot greater than 4"

suspected surgical site infection (redness, increasing pain, cloudy drainage, foul odor)

Recovery Objectives

0-6 weeks control pain, maintain some mobility through pendulum exercises while allowing the fracture to heal

>6 weeks start physical therapy and regain full range of motion

www.mccormickortho.com for more information more educational videos and postoperative protocols