

Reverse Shoulder Replacement—Dr. McCormick

What to Expect After Reverse Shoulder Replacement

Activity Restrictions

you may use your hand in front of your body for LIGHT tasks such as typing, writing or holding items about the same weight as a SMALL cup of coffee
you may drive when out of sling - you must be able to react quickly so try it in an empty parking lot if unsure

Surgical site care

keep your initial dressing dry
it is normal to notice some drainage/discoloration of the dressing
nurses at the hospital will change your dressing before you are discharged
you may shower with the island dressings on but do not scrub the area

Pain management

Rest wear the sling until the block wears off
remove the sling 3 times per day to:
move your hand, wrist and elbow to keep these from getting stiff
do shoulder pendulum exercises
the sutures holding your repaired tendons are not strong enough to withstand the force of normal use

Ice 30 minutes per hour – remember to keep something between the cold pack/ice and your skin to avoid frostbite

Take pain medications as prescribed. If you have a reaction to the medications then call the office at 304.343.4691

Follow-up clinic visit

usually at 2 weeks after surgery
we will check your surgical site, review Xrays and answer any questions you may have

When to call the office

reactions to medications
dressing discoloration spot greater than 4"
suspected surgical site infection (redness, increasing pain, cloudy drainage, foul odor)

Recovery Objectives

0-2 weeks control pain, maintain some mobility through pendulum exercises
2-6 weeks use arm IN FRONT OF BODY without sling
>6 weeks start physical therapy and regain full range of motion

www.mccormickortho.com for more information more educational videos and postoperative protocols